



# The impact of the Covid-19 pandemic on specialist palliative care services and how they have responded

**Prof Katherine Sleeman**


Laing Galazka Chair in Palliative Care  
King's College London  
Cicely Saunders Institute

Honorary Consultant in Palliative Medicine  
King's College Hospital NHS Trust

[www.csi.kcl.ac.uk](http://www.csi.kcl.ac.uk)

# Palliative care services have an essential front-line role... but early on this was not recognised

**Locally:** Sudden increase in palliative care team clinical activity from March 2020



**Nationally:** Essential front-line role of palliative care not recognised; palliative care community came together to raise awareness of needs and search for solutions



Urgent need for **research** to guide care and services

# March 2020: Rapid systematic review identified priorities for palliative care pandemic response

## SYSTEMS

### Policies

- **Maintain flexibility** and the ability to **rapidly modify or implement new policies**, such as changes to admission criteria.
- If visiting hours are limited, **consider** providing **daily telephone support to families**.

### Training and protocols

- **Provide protocols for non-specialists** on symptom management and provide basic palliative care training including communication
- **Consider** separate guidelines for **specific populations** such as those in care homes

### Communication and coordination

- **Identify** a **key decision maker** to improve communication, particularly if multiple professionals are involved outside of their usual practice
- **Involvement** in **triage systems** to assess likelihood of response to treatment and risk of dying
- **Share** protocols and advice within organisations

### Data

- **Standardise information collection & monitoring** to inform operational changes (for example number of patients seen, main symptoms and concerns, treatments, effectiveness of treatment, and outcomes)

## SPACE

### Maintain ability to move resources between settings

- **Consider moving resources** from inpatient to community setting according to need
- **Ensure** palliative care services are integrated into new systems for care, including community care centres and temporary hospitals

### Technology

- **Use virtual technology** to **enable communication** when visiting is restricted

## STAFF

### Deployment

- **Deploy staff flexibly** between palliative care settings according to need
- **Prioritise** sufficient **staff numbers**
- If patient contact with volunteer services needs to be restricted, **consider alternative roles** for example psychological support, bereavement support

### Skill mix

- **Involve chaplains and psychologists** with specialist palliative care expertise

### Resilience

- **Facilitate** camaraderie, **promote connectedness**, develop systems for reducing stress, and **support non-specialist staff**
- **Provide training in communication** and bereavement

## STUFF

### Medicines and equipment

- **Include** relevant **symptom medications in formularies** (to treat breathlessness, cough, fever, delirium, anxiety, agitation and pain)
- **Ensure** supplies of medications, **IV and subcutaneous** delivery systems and lines

### Personal protective equipment

- **Ensure** sufficient **supplies of PPE**

**Rapid systematic review** of role and response of hospice and palliative care services in epidemics

Etkind et al. 2020. The Role and Response of Palliative Care and Hospice Services in Epidemics and Pandemics: A Rapid Review to Inform Practice During the COVID-19 Pandemic. JPSM, 60 (1).



# CovPall study: Rapid evaluation of the Covid-19 pandemic response in PEOLC




**Survey** of hospice and palliative care services (Apr-Jul 2020); followed by study of symptoms & outcomes; and in depth case studies

458 responses: 277 UK, 85 Rest of Europe, 95 Rest of World


81% had cared for patients with suspected / confirmed COVID-19

Palliative care and hospice services were overwhelmed during the early pandemic but felt overlooked and lacked equipment, PPE, medicines and staff

a. Shortages by management type



b. Shortages by world regions



The  
Martin House  
RESEARCH CENTRE



International Observatory  
on End of Life Care




NIHR | Applied Research Collaboration  
South London



Medical  
Research  
Council



# Services felt overlooked, and not recognised as front line NHS




PPE shortages were common, often because services were not recognised as front line

*“Masks and gowns – as we are not classed as ‘frontline NHS’ we were unable to access from our local NHS board. .... this was less than satisfactory for 7 weeks, however this has now been resolved.” (Scotland)*

*“Tried to liaise with procurements services, NHS suppliers/CCG’s/local hospitals, appealing to local community through social media liaising with community groups and other organisations. Drawing on local networks available to us e.g. GP surgeries, schools, dentists, vets, local hardware companies, beauticians.” (South East)*

# Deaths shifted from hospice inpatient units into the community



## Population data for England and Wales

Number of deaths per week in care homes, hospitals and at home increased

Hospice deaths fell by 20% during first wave of Covid-19


# Hospice and palliative care services reported being more busy than before the pandemic

## CovPall data for UK services

48% (133) reported being **more busy**

Busyness was consequence of:

- Shifting resources in response to needs
- Telephone and remote working
- Educating, upskilling and supporting



Due to rounding, percentages may not total 100%.

*“One hospice unit has closed and the other has had to reduce beds [as they are] too close together. This has led to increase in community workload with more complex and unwell people at home requiring assessment, care coordination, clinical decision making and support.”* (Scotland)

*“We are supporting patients to remain in their usual place of care, home or care home. Patients/carers have been more psychologically distressed, and this is requiring more prolonged non face-to-face consultations.”* (North East)

*“We have also been engaging in much more education to healthcare professionals across all settings, including the care homes.”* (Wales)



# Hospice and palliative care services made rapid changes and 'frugal innovations'



Services made rapid changes within a context of ***crisis***

Services have been flexible, highly adaptive and have adopted a '**frugal innovation**' model

Hospices used, adjusted and recombined existing resources, structures and processes to deliver care.

## 'The crisis context'

### Barriers to change

- Fear and anxiety
- Duplication of effort
- Information overload
- IT infrastructure issues
- Funding issues

### Changes involved

- Streamlining access
- Extending services
- Increasing outreach
- Using communication technology
- Implementing innovations for staff wellbeing


### Enablers to change

- Pooling of staffing resources
- Staff flexibility
- Pre-existing IT infrastructure
- Collaborative teamwork
- Strong leadership

# Palliative care and hospice services experiences of advance care planning




277 UK hospice and palliative care services; 38% did more **direct** advance care planning; 59% provided more **support to others**



*‘...on the ground support in the care homes. This has been a collaborative approach...’*

[Mixed setting Palliative Care Team, Adult Service, Wales]

# CovPall-Rehab study: Impact on delivery of rehabilitation within palliative care services



Online survey, 61 rehabilitation services, 56 in hospices

Rapid adoption of digital and remote care delivery by rehabilitation teams changed capacity to provide and participate in rehabilitation

For some groups this led to enhanced access, teams could expand and reach new people (e.g., in rural areas, younger, unable to travel to the hospice building)

For others, there were concerns about a digital divide (e.g. with communication difficulties, no access to internet)

Hybrid models may expand the equity of access and reach of rehabilitation within specialist palliative care




# The response of services to people from ethnic minority groups



277 UK services; 34% had cared for people with Covid-19 from ethnic minority groups

Restricted visiting and communication challenges impacted disproportionately on ethnic minority groups

Findings identify need for **equitable** (not just equal) palliative care response



# CovPall-Connect study: Evaluation of the Covid response, connecting to boost impact and data assets



## News

### Health Data Research UK selects 12 projects to accelerate use of data for vital COVID-19 research

21 December 2020

12 urgent research projects are to receive new funding following a rapid call for COVID-19 data research initiatives by Health Data Research UK, Office for National Statistics and UK Research and...

HDRUK funded project

We will link CovPall data to other sources of nationally collected data (ONS mortality, HES data)

What is the relationship between palliative care responses to Covid-19 and outcomes such as admissions, discharges, place of death?



# Conclusions

- Palliative care and hospice services have had essential, front-line roles during Covid-19...
- ...but they felt overwhelmed during the early pandemic, overlooked, and lacked equipment, PPE, medicines and staff
- Care shifted from inpatient hospice settings into the community
- Hospice and palliative care services made rapid changes, adopting a 'frugal innovation' model
- Services provided direct and indirect care, collaborating across settings
- Disproportionate impact on ethnic minority groups, highlighting need for equitable care
- CovPall study highlights the power of clinical-academic collaborative research



# Acknowledgements

**CovPall Study Team:** Professor Irene J Higginson (Chief Investigator), Dr Sabrina Bajwah (Co-I), Dr Matthew Maddocks (Co-I), Professor Fliss Murtagh (Co-I) , Professor Nancy Preston (Co-I) , Dr Katherine E Sleeman (Co-I), Professor Catherine Walshe (Co-I), Professor Lorna K Fraser (Co-I), Dr Mevhibe B Hocaoglu (Co-I), Dr Adejoke Oluyase (Co-I), Dr Andrew Bradshaw, Lesley Dunleavy and Rachel L Cripps.

**CovPall Study Partners:** Hospice UK, Marie Curie, Sue Ryder, Palliative Outcome Scale Team, European Association of Palliative Care (EAPC), Together for Short Lives and Scottish Partnership for Palliative Care.

**PPI Steering Group Members:** Rashmi Kumar, Hameed Khan and Deb Smith

## Funding

This research was supported by Medical Research Council grant number MR/V012908/1. Additional support was from the National Institute for Health Research (NIHR), Applied Research Collaboration, South London, hosted at King's College Hospital NHS Foundation Trust, and Cicely Saunders International (Registered Charity No. 1087195).

**Contact me:**

[katherine.sleeman@kcl.ac.uk](mailto:katherine.sleeman@kcl.ac.uk)

@kesleeman